

Colombia - The probabilities of a hip fracture in women

The following tables give the 10-year probability (%) of a hip fracture according to the T-score for femoral neck BMD , the number of clinical risk factors (CRF) and age. Each table provides a mean estimate and a range, based on the epidemiology of Colombia. The range is not a confidence interval, but because the weight of different risk factors varies, is a true range.

Note that the BMI is set at 24 kg/m².

Age = 50 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	6.0	3.1	1.6	0.8	0.4	0.2	0.1	0.1	0.0	0.0	0.0
1	9.5 (6.3-12)	5.0 (3.3-6.5)	2.6 (1.7-3.4)	1.3 (0.9-1.7)	0.7 (0.4-0.9)	0.3 (0.2-0.5)	0.2 (0.1-0.2)	0.1 (0.1-0.1)	0.0 (0.0-0.1)	0.0 (0.0-0.0)	0.0 (0.0-0.0)
2	15 (8.9-22)	7.9 (4.6-12)	4.1 (2.4-6.2)	2.1 (1.2-3.2)	1.1 (0.6-1.7)	0.6 (0.3-0.9)	0.3 (0.2-0.4)	0.1 (0.1-0.2)	0.1 (0.0-0.1)	0.0 (0.0-0.1)	0.0 (0.0-0.0)
3	22 (13-34)	12 (7.0-19)	6.4 (3.6-10)	3.3 (1.9-5.5)	1.7 (1.0-2.8)	0.9 (0.5-1.5)	0.5 (0.3-0.8)	0.2 (0.1-0.4)	0.1 (0.1-0.2)	0.1 (0.0-0.1)	0.0 (0.0-0.1)
4	33 (22-47)	18 (12-28)	9.9 (6.2-16)	5.2 (3.2-8.3)	2.7 (1.6-4.3)	1.4 (0.8-2.2)	0.7 (0.4-1.1)	0.4 (0.2-0.6)	0.2 (0.1-0.3)	0.1 (0.1-0.2)	0.1 (0.0-0.1)
5	46 (36-59)	27 (21-37)	15 (11-21)	8.0 (5.9-12)	4.2 (3.0-6.1)	2.2 (1.6-3.1)	1.1 (0.8-1.6)	0.6 (0.4-0.8)	0.3 (0.2-0.5)	0.2 (0.1-0.2)	0.1 (0.1-0.1)
6	61	39	22	12	6.4	3.3	1.7	0.9	0.5	0.3	0.1

Ten year probability of hip fractures (%) according to BMD, the number of clinical risk factors (CRF) and age in women from Colombia.

Age = 55 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	7.7	4.2	2.3	1.2	0.7	0.4	0.2	0.1	0.1	0.0	0.0
1	12 (8.1-15)	6.6 (4.5-8.2)	3.6 (2.4-4.5)	2.0 (1.3-2.4)	1.1 (0.7-1.3)	0.6 (0.4-0.7)	0.3 (0.2-0.4)	0.2 (0.1-0.2)	0.1 (0.1-0.1)	0.1 (0.0-0.1)	0.0 (0.0-0.0)
2	18 (11-25)	10 (6.3-15)	5.7 (3.4-8.2)	3.1 (1.9-4.5)	1.7 (1.0-2.4)	0.9 (0.5-1.3)	0.5 (0.3-0.7)	0.3 (0.2-0.4)	0.2 (0.1-0.2)	0.1 (0.1-0.1)	0.0 (0.0-0.1)
3	27 (17-39)	16 (9.4-24)	8.7 (5.2-13)	4.8 (2.8-7.5)	2.6 (1.5-4.1)	1.4 (0.8-2.2)	0.8 (0.4-1.2)	0.4 (0.2-0.7)	0.2 (0.1-0.4)	0.1 (0.1-0.2)	0.1 (0.0-0.1)
4	38 (27-52)	23 (16-33)	13 (8.7-20)	7.3 (4.8-11)	4.0 (2.6-6.2)	2.2 (1.4-3.4)	1.2 (0.8-1.8)	0.7 (0.4-1.0)	0.4 (0.2-0.6)	0.2 (0.1-0.3)	0.1 (0.1-0.2)
5	51 (43-64)	33 (27-44)	19 (15-27)	11 (8.6-16)	6.1 (4.8-8.7)	3.3 (2.6-4.8)	1.8 (1.4-2.6)	1.0 (0.8-1.5)	0.6 (0.4-0.8)	0.3 (0.2-0.5)	0.2 (0.1-0.3)
6	66	45	28	16	9.1	5.0	2.7	1.5	0.9	0.5	0.3

Age = 60 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	8.4	4.8	2.7	1.5	0.9	0.5	0.3	0.2	0.1	0.1	0.0
1	13 (8.8-15)	7.4 (5.0-8.6)	4.2 (2.8-4.9)	2.4 (1.6-2.8)	1.3 (0.9-1.6)	0.7 (0.5-0.9)	0.4 (0.3-0.5)	0.2 (0.2-0.3)	0.1 (0.1-0.2)	0.1 (0.1-0.1)	0.0 (0.0-0.1)
2	19 (12-25)	11 (7.1-15)	6.4 (4.0-8.8)	3.6 (2.3-5.0)	2.0 (1.3-2.8)	1.1 (0.7-1.6)	0.6 (0.4-0.9)	0.4 (0.2-0.5)	0.2 (0.1-0.3)	0.1 (0.1-0.2)	0.1 (0.0-0.1)
3	27 (18-38)	17 (11-24)	9.7 (6.1-14)	5.5 (3.4-8.3)	3.1 (1.9-4.7)	1.8 (1.1-2.7)	1.0 (0.6-1.5)	0.6 (0.4-0.9)	0.3 (0.2-0.5)	0.2 (0.1-0.3)	0.1 (0.1-0.2)
4	38 (29-51)	24 (17-34)	14 (10-21)	8.3 (5.8-12)	4.7 (3.3-7.1)	2.7 (1.8-4.1)	1.5 (1.0-2.3)	0.9 (0.6-1.4)	0.5 (0.4-0.8)	0.3 (0.2-0.5)	0.2 (0.1-0.3)
5	51 (45-63)	34 (29-44)	21 (18-28)	12 (10-17)	7.1 (5.9-10)	4.0 (3.3-5.7)	2.3 (1.9-3.3)	1.3 (1.1-1.9)	0.8 (0.6-1.1)	0.5 (0.4-0.7)	0.3 (0.2-0.4)
6	65	46	29	18	10	6.0	3.5	2.0	1.2	0.7	0.4

Ten year probability of hip fractures (%) according to BMD, the number of clinical risk factors (CRF) and age in women from Colombia.

Age = 65 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	8.4	5.0	2.9	1.7	1.0	0.6	0.4	0.2	0.1	0.1	0.1
1	12 (8.8-15)	7.5 (5.2-8.8)	4.4 (3.1-5.3)	2.6 (1.8-3.1)	1.5 (1.0-1.8)	0.9 (0.6-1.1)	0.5 (0.4-0.6)	0.3 (0.2-0.4)	0.2 (0.1-0.2)	0.1 (0.1-0.1)	0.1 (0.1-0.1)
2	18 (12-23)	11 (7.3-14)	6.6 (4.3-8.6)	3.9 (2.5-5.1)	2.3 (1.5-3.0)	1.3 (0.9-1.8)	0.8 (0.5-1.1)	0.5 (0.3-0.6)	0.3 (0.2-0.4)	0.2 (0.1-0.2)	0.1 (0.1-0.1)
3	26 (18-35)	16 (11-22)	9.8 (6.5-14)	5.8 (3.9-8.4)	3.4 (2.3-5.0)	2.0 (1.3-2.9)	1.2 (0.8-1.8)	0.7 (0.5-1.1)	0.4 (0.3-0.7)	0.3 (0.2-0.4)	0.2 (0.1-0.2)
4	36 (28-47)	23 (18-32)	14 (11-20)	8.6 (6.4-12)	5.1 (3.8-7.5)	3.0 (2.2-4.4)	1.8 (1.3-2.7)	1.1 (0.8-1.6)	0.7 (0.5-1.0)	0.4 (0.3-0.6)	0.2 (0.2-0.4)
5	47 (42-58)	32 (28-41)	20 (17-27)	13 (10-17)	7.5 (6.2-10)	4.5 (3.7-6.2)	2.7 (2.2-3.8)	1.6 (1.3-2.3)	1.0 (0.8-1.4)	0.6 (0.5-0.9)	0.4 (0.3-0.5)
6	60	43	28	18	11	6.5	4.0	2.4	1.5	0.9	0.5

Age = 70 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	8.2	5.1	3.1	1.9	1.2	0.7	0.4	0.3	0.2	0.1	0.1
1	13 (11-17)	8.3 (7.1-11)	5.2 (4.4-6.5)	3.2 (2.7-4.0)	1.9 (1.6-2.5)	1.2 (1.0-1.5)	0.7 (0.6-1.0)	0.5 (0.4-0.6)	0.3 (0.3-0.4)	0.2 (0.2-0.2)	0.1 (0.1-0.2)
2	21 (17-27)	13 (11-18)	8.4 (6.6-11)	5.2 (4.1-7.1)	3.2 (2.5-4.4)	2.0 (1.5-2.7)	1.3 (1.0-1.7)	0.8 (0.6-1.1)	0.5 (0.4-0.7)	0.3 (0.2-0.4)	0.2 (0.2-0.3)
3	32 (25-39)	21 (16-27)	14 (10-17)	8.5 (6.2-11)	5.3 (3.8-7.0)	3.3 (2.4-4.4)	2.1 (1.5-2.8)	1.3 (0.9-1.8)	0.8 (0.6-1.1)	0.5 (0.4-0.7)	0.3 (0.2-0.5)
4	45 (36-52)	32 (24-37)	21 (16-25)	14 (10-17)	8.6 (6.2-11)	5.4 (3.9-6.7)	3.4 (2.5-4.3)	2.2 (1.6-2.7)	1.4 (1.0-1.7)	0.9 (0.6-1.1)	0.5 (0.4-0.7)
5	60 (52-64)	45 (38-50)	32 (26-36)	21 (17-24)	14 (11-16)	8.7 (6.8-10)	5.6 (4.3-6.5)	3.6 (2.8-4.1)	2.3 (1.8-2.6)	1.4 (1.1-1.7)	0.9 (0.7-1.1)
6	73	60	45	32	21	14	9.1	5.8	3.7	2.4	1.5

Ten year probability of hip fractures (%) according to BMD, the number of clinical risk factors (CRF) and age in women from Colombia.

Age = 75 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	7.5	4.8	3.1	2.0	1.3	0.8	0.5	0.4	0.2	0.2	0.1
1	14 (10-28)	9.3 (6.8-19)	6.1 (4.4-13)	3.9 (2.8-8.3)	2.5 (1.8-5.4)	1.7 (1.2-3.6)	1.1 (0.8-2.4)	0.7 (0.5-1.6)	0.5 (0.3-1.0)	0.3 (0.2-0.7)	0.2 (0.1-0.4)
2	24 (15-42)	16 (9.5-30)	11 (6.2-21)	7.3 (4.0-14)	4.7 (2.5-9.2)	3.1 (1.7-6.2)	2.1 (1.1-4.1)	1.4 (0.7-2.8)	0.9 (0.5-1.8)	0.6 (0.3-1.2)	0.4 (0.2-0.8)
3	37 (21-54)	27 (14-41)	19 (9.3-30)	13 (6.0-21)	8.4 (3.9-14)	5.6 (2.5-9.5)	3.8 (1.7-6.5)	2.5 (1.1-4.3)	1.7 (0.7-2.9)	1.1 (0.5-1.9)	0.7 (0.3-1.3)
4	53 (30-65)	41 (21-53)	30 (14-40)	21 (9.4-29)	14 (6.1-20)	9.8 (4.1-14)	6.6 (2.7-9.6)	4.5 (1.8-6.5)	3.0 (1.2-4.4)	2.0 (0.8-2.9)	1.3 (0.5-2.0)
5	68 (44-74)	57 (32-63)	45 (23-51)	33 (15-38)	23 (10-27)	16 (6.9-19)	11 (4.6-13)	7.6 (3.1-9.2)	5.1 (2.1-6.2)	3.4 (1.4-4.2)	2.3 (0.9-2.8)
6	79	72	61	48	36	26	18	13	8.7	5.9	3.9

Age = 80 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	6.3	4.3	2.9	1.9	1.3	0.9	0.6	0.4	0.3	0.2	0.1
1	12 (8.2-24)	8.0 (5.6-17)	5.5 (3.8-12)	3.7 (2.5-8.1)	2.6 (1.7-5.6)	1.8 (1.2-3.9)	1.2 (0.8-2.7)	0.9 (0.6-1.9)	0.6 (0.4-1.3)	0.4 (0.3-0.9)	0.3 (0.2-0.6)
2	20 (11-35)	14 (7.8-26)	9.8 (5.3-19)	6.7 (3.6-13)	4.7 (2.4-9.2)	3.3 (1.7-6.5)	2.3 (1.2-4.6)	1.6 (0.8-3.2)	1.1 (0.6-2.3)	0.8 (0.4-1.6)	0.5 (0.3-1.1)
3	30 (16-46)	23 (11-36)	16 (7.9-26)	12 (5.4-19)	8.1 (3.7-14)	5.8 (2.6-9.7)	4.1 (1.8-6.9)	2.9 (1.2-4.9)	2.0 (0.9-3.4)	1.4 (0.6-2.4)	1.0 (0.4-1.7)
4	43 (23-56)	34 (17-45)	26 (12-35)	19 (8.1-26)	13 (5.6-19)	9.7 (4.0-14)	6.9 (2.8-10)	4.9 (1.9-7.3)	3.4 (1.4-5.2)	2.4 (0.9-3.7)	1.7 (0.7-2.6)
5	57 (33-64)	47 (25-54)	38 (18-44)	28 (13-34)	21 (9.0-26)	16 (6.4-19)	11 (4.6-14)	8.1 (3.2-10)	5.8 (2.3-7.3)	4.1 (1.6-5.1)	2.9 (1.1-3.6)
6	70	61	51	41	32	24	18	13	9.4	6.7	4.7

Ten year probability of hip fractures (%) according to BMD, the number of clinical risk factors (CRF) and age in women from Colombia.

Age = 85 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	5.1	3.6	2.6	1.8	1.3	1.0	0.7	0.5	0.4	0.3	0.2
1	9.3 (6.6-20)	6.8 (4.7-15)	4.9 (3.4-11)	3.5 (2.4-7.7)	2.6 (1.8-5.7)	1.9 (1.3-4.2)	1.4 (0.9-3.1)	1.0 (0.7-2.3)	0.8 (0.5-1.7)	0.5 (0.4-1.2)	0.4 (0.3-0.9)
2	16 (9.2-28)	12 (6.6-21)	8.5 (4.8-16)	6.2 (3.4-12)	4.6 (2.5-8.9)	3.4 (1.8-6.7)	2.5 (1.3-5.0)	1.9 (1.0-3.7)	1.4 (0.7-2.7)	1.0 (0.5-2.0)	0.7 (0.4-1.5)
3	24 (13-37)	19 (9.3-30)	14 (6.8-23)	10 (4.9-17)	7.8 (3.6-13)	5.9 (2.7-9.9)	4.4 (2.0-7.5)	3.3 (1.5-5.6)	2.4 (1.1-4.1)	1.8 (0.8-3.1)	1.3 (0.6-2.3)
4	35 (18-46)	28 (14-38)	22 (10-30)	17 (7.3-23)	13 (5.4-18)	9.7 (4.1-14)	7.3 (3.0-10)	5.5 (2.2-7.9)	4.1 (1.6-5.9)	3.0 (1.2-4.4)	2.2 (0.9-3.3)
5	47 (25-53)	39 (20-45)	32 (15-37)	25 (11-30)	20 (8.3-24)	15 (6.3-18)	12 (4.7-14)	8.9 (3.5-11)	6.7 (2.6-8.3)	5.0 (1.9-6.2)	3.7 (1.4-4.7)
6	60	52	44	36	29	23	18	14	11	8.1	6.1

Age = 90 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	3.6	2.7	2.0	1.6	1.2	0.9	0.7	0.6	0.4	0.3	0.3
1	6.7 (4.8-14)	5.1 (3.6-11)	3.9 (2.7-8.5)	2.9 (2.0-6.5)	2.3 (1.6-5.1)	1.8 (1.2-4.0)	1.4 (1.0-3.1)	1.1 (0.7-2.4)	0.8 (0.6-1.9)	0.7 (0.4-1.5)	0.5 (0.3-1.1)
2	11 (6.5-21)	8.8 (5.0-16)	6.8 (3.8-13)	5.2 (2.9-9.9)	4.1 (2.2-7.9)	3.2 (1.7-6.2)	2.5 (1.4-4.9)	2.0 (1.1-3.8)	1.5 (0.8-3.0)	1.2 (0.6-2.3)	0.9 (0.5-1.8)
3	18 (9.1-29)	14 (6.9-23)	11 (5.3-18)	8.7 (4.0-14)	6.9 (3.2-12)	5.5 (2.5-9.3)	4.3 (1.9-7.3)	3.4 (1.5-5.8)	2.6 (1.2-4.5)	2.1 (0.9-3.5)	1.6 (0.7-2.8)
4	27 (13-37)	22 (10-30)	18 (7.8-24)	14 (6.1-20)	11 (4.8-16)	9.0 (3.8-13)	7.1 (2.9-10)	5.6 (2.3-8.0)	4.4 (1.8-6.3)	3.5 (1.4-5.0)	2.7 (1.1-3.9)
5	38 (19-44)	32 (15-37)	26 (12-31)	21 (9.0-25)	17 (7.2-21)	14 (5.7-17)	11 (4.5-14)	9.1 (3.5-11)	7.2 (2.8-8.8)	5.7 (2.2-7.0)	4.5 (1.7-5.5)
6	50	44	37	31	26	21	17	14	11	9.0	7.1