

Hungary - The probabilities of a hip fracture in men

The following tables give the 10-year probability (%) of a hip fracture according to the T-score for femoral neck BMD , the number of clinical risk factors (CRF) and age. Each table provides a mean estimate and a range, based on the epidemiology of Hungary. The range is not a confidence interval, but because the weight of different risk factors varies, is a true range.

Note that the BMI is set at 24 kg/m².

Age = 50 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	15	8.2	4.4	2.4	1.3	0.7	0.4	0.2	0.1	0.1	0.0
1	22 (16-29)	13 (8.6-17)	6.9 (4.6-9.2)	3.7 (2.5-5.0)	2.0 (1.3-2.7)	1.0 (0.7-1.4)	0.6 (0.4-0.7)	0.3 (0.2-0.4)	0.2 (0.1-0.2)	0.1 (0.1-0.1)	0.0 (0.0-0.1)
2	31 (21-43)	19 (12-27)	11 (6.5-15)	5.7 (3.5-8.6)	3.1 (1.9-4.6)	1.6 (1.0-2.5)	0.9 (0.5-1.3)	0.5 (0.3-0.7)	0.2 (0.2-0.4)	0.1 (0.1-0.2)	0.1 (0.0-0.1)
3	42 (30-55)	27 (18-37)	16 (9.8-23)	8.7 (5.3-13)	4.7 (2.8-7.2)	2.5 (1.5-3.9)	1.4 (0.8-2.1)	0.7 (0.4-1.1)	0.4 (0.2-0.6)	0.2 (0.1-0.3)	0.1 (0.1-0.2)
4	54 (42-65)	37 (26-49)	23 (15-32)	13 (8.4-19)	7.2 (4.5-11)	3.9 (2.4-5.9)	2.1 (1.3-3.2)	1.1 (0.7-1.7)	0.6 (0.4-0.9)	0.3 (0.2-0.5)	0.2 (0.1-0.3)
5	65 (56-73)	48 (39-58)	31 (24-41)	19 (14-26)	11 (7.7-15)	5.8 (4.2-8.3)	3.1 (2.2-4.5)	1.7 (1.2-2.4)	0.9 (0.6-1.3)	0.5 (0.4-0.7)	0.3 (0.2-0.4)
6	74	60	42	27	16	8.7	4.7	2.5	1.4	0.8	0.4

Ten year probability of hip fractures (%) according to BMD, the number of clinical risk factors (CRF) and age in men from Hungary.

Age = 55 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	16	9.3	5.2	2.9	1.6	0.9	0.5	0.3	0.2	0.1	0.1
1	23 (17-29)	14 (9.7-17)	7.9 (5.5-10)	4.5 (3.1-5.7)	2.5 (1.7-3.2)	1.4 (0.9-1.8)	0.8 (0.5-1.0)	0.4 (0.3-0.5)	0.2 (0.2-0.3)	0.1 (0.1-0.2)	0.1 (0.1-0.1)
2	32 (23-42)	20 (13-27)	12 (7.7-16)	6.7 (4.3-9.5)	3.8 (2.4-5.4)	2.1 (1.3-3.0)	1.2 (0.7-1.7)	0.6 (0.4-0.9)	0.4 (0.2-0.5)	0.2 (0.1-0.3)	0.1 (0.1-0.2)
3	42 (32-54)	28 (20-37)	17 (11-24)	10 (6.5-14)	5.7 (3.6-8.2)	3.2 (2.0-4.6)	1.8 (1.1-2.6)	1.0 (0.6-1.5)	0.6 (0.3-0.8)	0.3 (0.2-0.5)	0.2 (0.1-0.3)
4	53 (43-63)	37 (28-47)	24 (17-32)	14 (10-20)	8.4 (5.7-12)	4.8 (3.2-7.0)	2.7 (1.8-3.9)	1.5 (1.0-2.2)	0.8 (0.6-1.3)	0.5 (0.3-0.7)	0.3 (0.2-0.4)
5	63 (55-70)	48 (40-57)	33 (26-41)	21 (16-27)	12 (9.4-17)	7.0 (5.3-9.7)	4.0 (3.0-5.5)	2.2 (1.7-3.1)	1.3 (1.0-1.8)	0.7 (0.5-1.0)	0.4 (0.3-0.6)
6	71	58	43	28	17	10	5.8	3.3	1.9	1.1	0.6

Age = 60 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	14	8.3	5.0	2.9	1.7	1.0	0.6	0.3	0.2	0.1	0.1
1	19 (14-23)	12 (8.7-14)	7.4 (5.2-8.8)	4.4 (3.1-5.3)	2.6 (1.8-3.1)	1.5 (1.1-1.8)	0.9 (0.6-1.1)	0.5 (0.4-0.6)	0.3 (0.2-0.4)	0.2 (0.1-0.2)	0.1 (0.1-0.1)
2	27 (19-34)	17 (12-23)	11 (7.3-14)	6.5 (4.4-8.7)	3.9 (2.6-5.2)	2.3 (1.5-3.1)	1.3 (0.9-1.8)	0.8 (0.5-1.1)	0.5 (0.3-0.7)	0.3 (0.2-0.4)	0.2 (0.1-0.2)
3	36 (27-45)	24 (18-31)	15 (11-21)	9.4 (6.6-13)	5.7 (3.9-7.8)	3.4 (2.3-4.7)	2.0 (1.3-2.8)	1.2 (0.8-1.7)	0.7 (0.5-1.0)	0.4 (0.3-0.6)	0.3 (0.2-0.4)
4	45 (37-55)	32 (25-40)	21 (16-28)	13 (9.9-18)	8.2 (6.0-11)	5.0 (3.6-7.0)	3.0 (2.1-4.2)	1.7 (1.2-2.5)	1.1 (0.8-1.5)	0.6 (0.5-0.9)	0.4 (0.3-0.6)
5	55 (49-63)	41 (36-50)	29 (24-36)	19 (15-25)	12 (9.6-16)	7.2 (5.8-9.7)	4.3 (3.5-5.9)	2.6 (2.1-3.5)	1.6 (1.3-2.2)	1.0 (0.8-1.3)	0.6 (0.5-0.8)
6	64	51	37	26	16	10	6.2	3.7	2.3	1.4	0.9

Ten year probability of hip fractures (%) according to BMD, the number of clinical risk factors (CRF) and age in men from Hungary.

Age = 65 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	11	7.0	4.4	2.8	1.7	1.1	0.7	0.4	0.3	0.2	0.1
1	15 (11-17)	9.9 (7.3-11)	6.4 (4.6-7.2)	4.0 (2.9-4.6)	2.5 (1.8-2.9)	1.6 (1.1-1.8)	1.0 (0.7-1.1)	0.6 (0.4-0.7)	0.4 (0.3-0.5)	0.3 (0.2-0.3)	0.2 (0.1-0.2)
2	21 (16-26)	14 (10-17)	9.1 (6.5-11)	5.8 (4.1-7.4)	3.7 (2.6-4.7)	2.3 (1.6-3.0)	1.4 (1.0-1.9)	0.9 (0.6-1.2)	0.6 (0.4-0.8)	0.4 (0.3-0.5)	0.2 (0.2-0.3)
3	28 (21-35)	19 (14-25)	13 (9.4-17)	8.2 (6.0-11)	5.3 (3.8-7.0)	3.3 (2.4-4.5)	2.1 (1.5-2.8)	1.3 (1.0-1.8)	0.9 (0.6-1.2)	0.5 (0.4-0.7)	0.4 (0.3-0.5)
4	36 (30-44)	25 (21-32)	17 (14-22)	12 (9.0-15)	7.5 (5.8-10)	4.8 (3.6-6.4)	3.0 (2.3-4.1)	1.9 (1.5-2.6)	1.2 (0.9-1.7)	0.8 (0.6-1.1)	0.5 (0.4-0.7)
5	44 (40-51)	33 (29-40)	23 (20-29)	16 (14-20)	10 (9.0-14)	6.7 (5.8-9.0)	4.3 (3.7-5.8)	2.7 (2.3-3.7)	1.8 (1.5-2.4)	1.2 (1.0-1.6)	0.7 (0.6-1.0)
6	52	41	30	21	14	9.4	6.1	3.9	2.5	1.7	1.1

Age = 70 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	9.5	6.5	4.4	2.9	1.9	1.3	0.9	0.6	0.4	0.3	0.2
1	14 (13-19)	10 (8.8-13)	6.8 (6.0-9.0)	4.6 (4.1-6.1)	3.1 (2.7-4.1)	2.1 (1.8-2.7)	1.4 (1.2-1.8)	0.9 (0.8-1.2)	0.6 (0.6-0.8)	0.4 (0.4-0.6)	0.3 (0.3-0.4)
2	21 (17-26)	15 (12-19)	10 (8.4-13)	7.2 (5.7-9.2)	4.8 (3.9-6.3)	3.2 (2.6-4.3)	2.2 (1.7-2.9)	1.5 (1.2-1.9)	1.0 (0.8-1.3)	0.7 (0.6-0.9)	0.5 (0.4-0.6)
3	29 (23-36)	22 (17-27)	16 (12-19)	11 (8.4-14)	7.5 (5.7-9.5)	5.1 (3.9-6.5)	3.4 (2.6-4.4)	2.3 (1.8-3.0)	1.6 (1.2-2.1)	1.1 (0.8-1.4)	0.7 (0.6-1.0)
4	39 (31-45)	30 (24-36)	22 (17-27)	16 (12-20)	11 (8.5-14)	7.8 (5.8-9.6)	5.3 (4.0-6.5)	3.6 (2.7-4.5)	2.5 (1.9-3.1)	1.7 (1.3-2.1)	1.2 (0.9-1.5)
5	49 (41-53)	40 (32-44)	31 (24-34)	23 (18-26)	17 (13-19)	12 (8.8-13)	8.1 (6.0-9.1)	5.6 (4.2-6.3)	3.9 (2.9-4.4)	2.7 (2.0-3.0)	1.9 (1.4-2.1)
6	59	50	40	32	24	17	12	8.6	6.1	4.2	2.9

Ten year probability of hip fractures (%) according to BMD, the number of clinical risk factors (CRF) and age in men from Hungary.

Age = 75 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	10	7.3	5.3	3.8	2.7	1.9	1.3	1.0	0.7	0.5	0.4
1	17 (12-32)	13 (9.3-26)	9.4 (6.8-20)	6.9 (4.9-15)	5.0 (3.5-11)	3.5 (2.5-7.8)	2.5 (1.8-5.6)	1.8 (1.3-4.1)	1.3 (1.0-3.0)	1.0 (0.7-2.2)	0.7 (0.5-1.6)
2	25 (16-41)	20 (12-34)	15 (9.2-27)	11 (6.7-21)	8.5 (4.9-16)	6.2 (3.5-11)	4.4 (2.5-8.3)	3.3 (1.8-6.1)	2.4 (1.3-4.5)	1.7 (1.0-3.3)	1.3 (0.7-2.4)
3	35 (21-49)	29 (16-42)	23 (12-34)	18 (9.3-27)	14 (6.8-21)	10 (4.9-16)	7.4 (3.6-12)	5.5 (2.6-8.9)	4.1 (1.9-6.7)	3.0 (1.4-4.9)	2.2 (1.0-3.6)
4	45 (27-57)	39 (22-50)	32 (17-42)	26 (13-34)	20 (9.4-27)	16 (6.9-21)	12 (5.0-16)	8.8 (3.7-12)	6.6 (2.8-9.3)	4.9 (2.0-6.9)	3.6 (1.5-5.1)
5	56 (36-63)	49 (29-56)	42 (23-49)	35 (18-42)	29 (14-34)	23 (10-27)	18 (7.4-21)	14 (5.6-17)	10 (4.1-13)	7.9 (3.1-9.5)	5.9 (2.3-7.1)
6	66	60	53	46	38	32	25	20	16	12	9.1

Age = 80 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	11	8.1	6.1	4.6	3.5	2.6	1.9	1.5	1.1	0.9	0.7
1	17 (13-34)	14 (10-28)	11 (7.8-22)	8.2 (5.9-17)	6.3 (4.5-14)	4.7 (3.4-10)	3.6 (2.5-8.0)	2.8 (1.9-6.2)	2.1 (1.5-4.8)	1.6 (1.1-3.7)	1.2 (0.9-2.8)
2	26 (17-43)	21 (13-36)	17 (10-30)	13 (7.7-24)	10 (5.8-19)	8.0 (4.4-15)	6.1 (3.3-12)	4.8 (2.6-9.1)	3.7 (2.0-7.1)	2.9 (1.5-5.5)	2.2 (1.2-4.2)
3	36 (22-51)	30 (17-44)	25 (14-38)	20 (10-31)	16 (8.1-26)	13 (6.1-20)	9.9 (4.7-16)	7.8 (3.6-13)	6.1 (2.8-10)	4.8 (2.1-8.0)	3.7 (1.6-6.2)
4	47 (28-58)	40 (23-52)	35 (18-45)	29 (14-39)	24 (11-32)	19 (8.5-27)	15 (6.5-21)	12 (5.1-17)	9.7 (4.0-14)	7.6 (3.1-11)	6.0 (2.4-8.6)
5	57 (36-63)	51 (30-58)	45 (25-52)	39 (20-45)	33 (16-38)	27 (12-32)	22 (9.6-26)	18 (7.6-22)	15 (5.9-18)	12 (4.6-14)	9.3 (3.6-11)
6	66	62	56	50	43	37	31	26	21	17	14

Ten year probability of hip fractures (%) according to BMD, the number of clinical risk factors (CRF) and age in men from Hungary.

Age = 85 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	8.9	7.0	5.5	4.4	3.4	2.7	2.1	1.7	1.4	1.1	0.9
1	15 (11-30)	12 (9.0-25)	9.8 (7.2-21)	7.8 (5.7-17)	6.2 (4.5-13)	4.9 (3.5-11)	3.9 (2.8-8.7)	3.2 (2.2-7.0)	2.5 (1.8-5.7)	2.0 (1.4-4.6)	1.6 (1.1-3.7)
2	23 (14-39)	19 (12-34)	16 (9.2-28)	13 (7.4-23)	10 (5.8-19)	8.3 (4.6-16)	6.7 (3.7-13)	5.5 (3.0-10)	4.4 (2.4-8.5)	3.6 (1.9-6.9)	2.9 (1.5-5.6)
3	33 (19-47)	28 (16-42)	24 (13-36)	20 (10-31)	16 (8.1-26)	13 (6.4-21)	11 (5.1-18)	8.9 (4.1-15)	7.3 (3.4-12)	6.0 (2.7-10)	4.8 (2.2-8.1)
4	44 (25-55)	38 (21-50)	33 (17-44)	29 (14-38)	24 (11-33)	20 (9.0-28)	17 (7.3-23)	14 (5.9-20)	12 (4.8-16)	9.5 (3.9-14)	7.8 (3.2-11)
5	54 (34-61)	49 (29-56)	44 (24-50)	39 (20-45)	34 (16-39)	29 (13-34)	24 (11-29)	21 (8.8-24)	17 (7.2-21)	15 (5.9-17)	12 (4.8-14)
6	64	60	55	50	45	39	34	29	25	21	18

Age = 90 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	5.7	4.7	3.9	3.1	2.6	2.1	1.7	1.5	1.2	1.0	0.9
1	10 (7.4-21)	8.4 (6.1-18)	7.0 (5.0-15)	5.8 (4.1-13)	4.7 (3.4-10)	3.9 (2.7-8.6)	3.3 (2.3-7.3)	2.7 (1.9-6.2)	2.3 (1.6-5.2)	1.9 (1.4-4.4)	1.6 (1.1-3.7)
2	16 (9.6-29)	14 (8.0-25)	12 (6.6-21)	9.7 (5.4-18)	8.1 (4.5-15)	6.7 (3.7-13)	5.7 (3.1-11)	4.8 (2.6-9.1)	4.1 (2.2-7.7)	3.4 (1.8-6.6)	2.9 (1.5-5.5)
3	25 (13-37)	21 (11-33)	18 (9.1-28)	15 (7.6-25)	13 (6.2-21)	11 (5.1-18)	9.3 (4.3-15)	7.9 (3.6-13)	6.7 (3.1-11)	5.7 (2.6-9.5)	4.8 (2.2-8.1)
4	34 (18-45)	30 (15-41)	27 (13-36)	23 (11-32)	20 (8.9-27)	17 (7.4-23)	14 (6.2-20)	12 (5.3-18)	11 (4.5-15)	9.1 (3.8-13)	7.8 (3.2-11)
5	45 (25-52)	41 (21-47)	37 (18-42)	33 (15-38)	28 (13-33)	25 (11-29)	22 (9.2-25)	19 (7.8-22)	16 (6.7-19)	14 (5.7-17)	12 (4.8-14)
6	56	52	48	43	39	34	31	27	24	21	18